



**Live freeze-dried vaccine,
LIBDV strain**

**For the active immunization of
chickens against Infectious Bursal Disease**

► COMPOSITION

CEVAC® GUMBO L contains the LIBDV intermediate strain of Infectious Bursal Disease virus in live, freeze-dried form.

► INDICATIONS

For the active immunization of healthy chickens against Infectious Bursal Disease (Gumboro Disease).

► ADMINISTRATION AND DOSAGE

CEVAC® GUMBO L must be administered **by drinking water method**.

Broilers should be vaccinated with **CEVAC® GUMBO L** from 14 to 21 days of age, depending on the level of maternally derived antibodies. In case of heterogeneity of this level between chicks, a first vaccination between 7 to 10 days of age is recommended.

Pullets should be vaccinated twice between the age of 20 and 28 days, allowing a six-day interval between administrations. In case of heterogeneity of the maternally derived antibody level, a first vaccination between the age of 14 to 16 days is recommended.

The exact date of vaccination can be determined by checking the level of maternally derived antibodies.

► STORAGE

- Store vaccine between +2°C and +8°C or 35°F and 45°F
- Protect from light

► PACKAGED

- 1,000 - 2,500 and 5,000 dose vials
- 20 x 1,000 dose vials / box
- 20 x 2,500 dose vials / box
- 20 x 5,000 dose vials / box

Infectious Bursal Disease (Gumboro Disease)
LIVE VACCINES