



Inactivated oil emulsion AIV H9N2 strain vaccine

For the immunization of chickens against Avian Influenza subtype H9

► COMPOSITION

CEVAC® FLU H9 K contains subtype H9N2 (Middle East isolate) Avian Influenza virus in inactivated form in oily adjuvant.

► INDICATIONS

CEVAC® FLU H9 K is recommended in the preventive vaccination of broilers and layers or breeders pullets against Avian Influenza (H9N2).

► ADMINISTRATION AND DOSAGE

CEVAC® FLU H9 K should be injected by subcutaneous or intramuscular route to:

- **Broilers** at day old or at 7 days of age at 0.2 ml dose per bird.
- **Layers or breeders pullets** at 3-5 weeks of age at 0.2 ml dose per bird.
For pullets at 16-18 weeks of age inject 0.5 ml dose per bird, before the onset of lay.

► STORAGE

- Store vaccine between +2°C and +8°C or 35°F and 45°F
- Protect from light
- Do not freeze

► PACKAGED

Bottles of 500 ml:

- **Broilers**
at 0.2 ml dose / 2500 doses/bottle.
- **Layers or breeders pullets**
at 0.5 ml dose / 1000 doses/bottle.

WARNING

This product contains oil based adjuvant; accidental injection into a human may cause a serious localized reaction. Seek expert medical attention immediately.