



Inactivated oil emulsion vaccine

**For the immunization of chickens
against Newcastle Disease (ND)
and Infectious Bronchitis (IB)**



► COMPOSITION

CEVAC® ND IB K contains a combination of the La Sota strain of Newcastle Disease virus and the Massachusetts strain of Infectious Bronchitis virus in inactivated form, homogenised with oil adjuvant.

► INDICATIONS

CEVAC® ND IB K is recommended for the vaccination of breeder and layingtype chicken flocks, previously immunised against Infectious Bronchitis (IB) and Newcastle Disease (ND) with attenuated live virus vaccines.

► ADMINISTRATION AND DOSAGE

CEVAC® ND IB K should be administered to breeder and laying-type pullets 3 to 4 weeks before the onset of lay, between 16 and 20 weeks of age.

The vaccine must be **injected subcutaneously** (under the skin of the neck) or by intramuscular route (breast or thigh muscles) at the dose of 0.5 ml per bird.

► STORAGE

- Store vaccine between +2°C and +8°C or 35°F and 45°F
- Protect from light
- Do not freeze

► PACKAGED

1,000 doses / 500 ml
5 x 1,000 dose bottles /box

WARNING

This product contains oil based adjuvant; accidental injection into a human may cause a serious localized reaction. Seek expert medical attention immediately.