



**Live freeze-dried vaccine,  
Massachusetts B48 strain**

**For the active immunization of  
chickens against Infectious Bronchitis**

### ► COMPOSITION

**CEVAC® MASS L** contains the Massachusetts B48 strain of Infectious Bronchitis virus in live, freeze-dried form. The embryonated hen eggs used in the production of the vaccine are obtained from specified-pathogen-free (SPF) flocks.

### ► INDICATIONS

For the active immunization of healthy chickens against Infectious Bronchitis. The vaccine may be used for initial or revaccination of broilers, pullets and hens.

### ► ADMINISTRATION AND DOSAGE

Primary vaccination should be carried-out with **CEVAC® MASS L** by **eye-drop or spray method**, as from 1 to 4 days of age.

Booster vaccination is recommended three to four weeks later by spray or drinking water.

For pullets, a second booster vaccination is recommended at 10 to 12 weeks of age.

### ► SIDE EFFECT

Rarely, catarrhal respiratory symptoms appear 4 to 6 days following vaccination but disappear in a few days.

### ► STORAGE

- Store vaccine between +2°C and +8°C or 35°F and 45°F

- Protect from light

### ► PACKAGED

1,000 and 2,500 dose vials

20 x 1,000 dose vials / box

20 x 2,500 dose vials / box