



Inactivated vaccine for the active immunization of chickens against Infectious Bronchitis (IB), Newcastle Disease (ND), Egg Drop Syndrome virus (EDS), Avian Metapneumovirus (AMP)



COMPOSITION

CEVAC® MEGAMUNE K is an inactivated multivalent vaccine recommended for the vaccination of layer and breeder type chicken flocks.

VACCINE TYPE & STRAIN

Inactivated oil emulsion vaccine:

- Newcastle disease virus strain NDV-S7 LaSota
- Avian Infectious Bronchitis virus strain M-41
- Avian Infectious Bronchitis virus strain OX Fr
- Egg Drop Syndrome virus strain B8/78
- Avian Metapneumovirus strain TRT50

INDICATIONS

CEVAC MEGAMUNE K is recommended for the vaccination of layer and breeder type chicken flocks, in order - to reduce mortality, clinical signs, lesions and virus shedding caused by Newcastle disease virus; - to reduce egg drop, respiratory signs and virus shedding caused by the Massachusetts and QX-like serotypes of Infectious bronchitis virus; - to reduce egg drop caused by the Egg drop syndrome virus; - to reduce respiratory signs (e.g. signs linked to Swollen Head Syndrome) and virus shedding caused by Avian metapneumovirus.es.

ADMINISTRATION AND DOSAGE

The dose of the vaccine is 0.5 ml, injected by intramuscular route (into breast muscles) or subcutaneously (into the back of the neck) from 14 weeks of age, no later than 3 weeks before expected onset of the laying period.

STORAGE

- Store and transport vaccine between +2°C and +8°C
- Protected from light
- Do not freeze

PACKAGED

1.000 doses / 500 ml

WARNING

This product contains oil based adjuvant; accidental injection into a human may cause a serious localized reaction.

Seek expert medical attention immediately.





